

# Semiclassical limit of the lowest eigenvalue of Schrödinger operators on path spaces

Shigeki Aida  
Osaka University

Let  $(X, g)$  be a compact Riemannian manifold and  $E, V$  be smooth real-valued functions on  $X$ . Let  $\mu_\lambda$  be the probability measure such that  $d\mu_\lambda = Z_\lambda^{-1} e^{-\lambda E} dx$  on  $X$ , where  $dx$  is the volume element and  $Z_\lambda$  is the normalizing constant.

Let

$$\mathcal{E}_\lambda(f, f) = \int_X |\nabla f(x)|^2 d\mu_\lambda(x) = \int_X (-L_\lambda f)(x) \cdot f(x) d\mu_\lambda(x) \quad (1)$$

$$\mathcal{E}_{\lambda, V}(f, f) = \mathcal{E}_\lambda(f, f) + \lambda^2 \int_X V(x) f(x)^2 d\mu_\lambda(x) \quad (2)$$

$$= \int_X (-L_{\lambda, V} f)(x) \cdot f(x) d\mu_\lambda(x) \quad (3)$$

$$E_0(\lambda, V) = \inf \left\{ \mathcal{E}_{\lambda, V}(f, f) \mid \|f\|_{L^2(\mu_\lambda)} = 1 \right\}. \quad (4)$$

We recall the asymptotic behavior of  $E_0(\lambda, V)$  as  $\lambda \rightarrow \infty$ . Assume that

(A1)  $U(x) = \frac{|\nabla E(x)|^2}{4} + V(x)$  is a nonnegative function which has finite zero point set  $\{c_1, \dots, c_n\}$ .

(A2) The Hessian of  $U$  at  $c_i$  ( $1 \leq i \leq n$ ) is strictly positive.

Then we have

$$\lim_{\lambda \rightarrow \infty} \frac{E_0(\lambda, V)}{\lambda} = \min_{1 \leq i \leq n} \operatorname{tr} \left\{ \sqrt{\frac{(\nabla^2 U)(c_i)}{2}} - \frac{\nabla^2 E}{2}(c_i) \right\}, \quad (5)$$

where  $\nabla^2$  denotes the second covariant derivative which is defined by the Levi-Civita connection. We extend this result to path space  $X$  with a Dirichlet form and a probability measure  $\mu_\lambda$  which is written formally as  $d\mu_\lambda(\gamma) = Z_\lambda^{-1} \exp(-\lambda E(\gamma)) d\gamma$ , where  $\gamma$  denotes a path and  $E(\gamma)$  is the energy of the path.  $d\gamma$  is a formal Riemannian volume. In this talk, we consider the following three cases.

(I)  $X$  is an abstract Wiener space  $(B, H, \mu)$ . The Dirichlet form is given by

$$\mathcal{E}_{\lambda, A}(f, f) = \int_B |A(w) Df(w)|_H^2 d\mu_\lambda(w). \quad (6)$$

Here  $D$  denotes the usual  $H$ -derivative,  $A(w) \in L(H, H)$  and  $\mu_\lambda(\cdot) = \mu(\sqrt{\lambda} \cdot)$ . We consider the Schrödinger operator  $-L_{\lambda, A, V}$  corresponding to a semi-bounded form  $\mathcal{E}_{\lambda, A, V}(f, f) = \mathcal{E}_{\lambda, A}(f, f) + \lambda^2 \int_B V(w) f(w)^2 d\mu_\lambda(w)$ . In this case,  $E(w) = \frac{1}{2} \|w\|_H^2$ . If  $A(w) = I_H$ , then the limit  $\lim_{\lambda \rightarrow \infty} \frac{E_0(\lambda, V)}{\lambda}$  was studied in [1].

(II)  $X = P_x(M) = C([0, 1] \rightarrow M \mid \gamma(0) = x)$ . Here  $M$  is a compact Riemannian manifold. The measure  $\mu_\lambda$  is the Brownian motion measure which is given by the heat semigroup  $e^{\frac{t}{2\lambda}\Delta}$ . The Dirichlet form is given by the  $H$ -derivative:

$$(\nabla F)(\gamma) = \sum_{i=1}^n \tau(\gamma)_{t_i}^{-1} (\nabla f)_{\gamma(t_i)}(\gamma(t_1), \dots, \gamma(t_n)) t \wedge t_i, \quad (7)$$

where  $F(\gamma) = f(\gamma(t_1), \dots, \gamma(t_n))$  and  $\tau(\gamma)_t : T_x(M) \rightarrow T_{\gamma(t)}M$  denotes the stochastic parallel translation. In this case,  $E(\gamma) = \frac{1}{2} \int_0^1 |\dot{\gamma}(t)|^2 dt$ .

(III)  $X = P_{e,a}(G) = C([0, 1] \rightarrow G \mid \gamma(0) = e, \gamma(1) = a \in G)$ . Here  $G$  is a compact Lie group,  $e$  is the unit element. In this case,  $\mu_\lambda$  is the pinned Brownian motion measure  $\nu_{\lambda,e,a}$  which is defined by  $e^{\frac{t}{2\lambda}\Delta}$ . The Dirichlet form is given by the probability measure  $\nu_{\lambda,e,a}$  and the  $H$ -derivative:

$$(\nabla f(\gamma), h) = \lim_{\varepsilon \rightarrow 0} \frac{f(e^{\varepsilon h(\cdot)} \gamma(\cdot)) - f(\gamma)}{\varepsilon}, \quad (8)$$

where  $h \in H^1([0, 1] \rightarrow \mathfrak{g} \mid h(0) = h(1) = 0)$  and  $\mathfrak{g}$  is the Lie algebra of  $G$ . In this case again,  $E(\gamma) = \frac{1}{2} \int_0^1 |\dot{\gamma}(t)|^2 dt$ .

In these cases, we can determine the limit  $\lim_{\lambda \rightarrow \infty} \frac{E_0(\lambda, V)}{\lambda}$  under the similar assumptions in (A1) and (A2). In the proof of the lower bound estimate, we combine the following two results:

- (1) Rough lower bound estimate on  $E_0(\lambda, V)$  which is given by a log-Sobolev inequality on  $X$
- (2) "Approximation" of the Schrödinger operator near zero points of the potential function  $U$  by a Schrödinger operator with a quadratic potential function on an abstract Wiener space in the case where  $A(w) = I_H + T$  and  $T$  is a trace class operator which is independent of  $w$ .

In the case of (II)( $X = P_x(M)$ ), as to (2), we use an infinite dimensional version of the following simple result and a pointwise estimate on  $\frac{L_\lambda \Omega_{\lambda, V}(x) - L_{\lambda, A} \Omega_{\lambda, V}(x)}{\Omega_{\lambda, V}(x)}$ .

**Proposition 1** *We consider the forms in (1) and (2) and the lowest eigenvalue  $E_0(\lambda, V)$  on the compact Riemannian manifold  $(X, g)$ . Let  $A(x) \in L(T_x X, T_x X)$  and assume that  $x \rightarrow A(x)$  is smooth and set*

$$\mathcal{E}_{\lambda, A}(f, f) = \int_X |A(x) \nabla f(x)|^2 d\mu_\lambda(x) = \int_X (-L_{\lambda, A} f)(x) \cdot f(x) d\mu_\lambda(x) \quad (9)$$

$$\mathcal{E}_{\lambda, A, V}(f, f) = \mathcal{E}_{\lambda, A}(f, f) + \lambda^2 \int_X V(x) f(x)^2 d\mu_\lambda(x). \quad (10)$$

Let  $\Omega_{\lambda, V}$  be the positive normalized eigenfunction of  $-L_{\lambda, V}$  corresponding to  $E_0(\lambda, V)$ . Then for any  $f \in C^\infty(X)$ , we have

$$\begin{aligned} \mathcal{E}_{\lambda, A}(f \Omega_{\lambda, V}, f \Omega_{\lambda, V}) &= \int_X |A(x) \nabla f(x)|^2 \Omega_{\lambda, V}(x)^2 d\mu_\lambda(x) + E_0(\lambda, V) \|f \Omega_{\lambda, V}\|_{L^2(\mu_\lambda)}^2 \\ &\quad + \int_X \frac{L_\lambda \Omega_{\lambda, V}(x) - L_{\lambda, A} \Omega_{\lambda, V}(x)}{\Omega_{\lambda, V}(x)} f(x)^2 \Omega_{\lambda, V}(x)^2 d\mu_\lambda(x). \end{aligned} \quad (11)$$

## References

- [1] S. Aida, Semiclassical limit of the lowest eigenvalue of a Schrödinger operator on a Wiener space, *J. Funct. Anal.* **203**, (2003), 401–424.